


Status of the XIS Orbital Calibration and Softwares

K. Hayashida, N. Tawa, M. Nagai, H. Nakajima, M. Uchino,
N. Anabuki, S. Katsuda, H. Uchida, H. Tsunemi (Osaka Univ.),
H. Matsumoto, H. Mori, H. Uchiyama, M. Ozawa, T. G. Tsuru,
Y. Hyodo, M. Nobukawa, K. Koyama (Kyoto Univ.),
T. Dotani, H. Murakami, A. Bamba, M. Ozaki, T. Anada
(JAXA),
D. Takei, S. Kitamoto (Rikkyo Univ.),
T. Kato, S. Maeno, K. Mori (Miyazaki Univ.),
Y. Ishisaki (TMU), T. Kohmura (Kohgakuinn Univ.),
E. Miller, B. LaMarr, S. E. Kissel, **M. W. Bautz** (MIT),
and the Suzaku Team


Current Status of the XIS

- No significant change in the XIS hardware since the last Users' Meeting 2007May
- SCI-ON for all the observations

Peak ch of Mn $K\alpha$ from ^{55}Fe calibration sources / that at the launch




FWHM of Mn $K\alpha$ from ^{55}Fe calibration sources




Ver. 2 Processing & HEASOFT6.4

- Ver. 2 processing started.
 - Full processing of the SCI-ON data
- HEASOFT v6.4
 - Updates + New software xisnxbgen is introduced
- CALDB updated (as always)


Quality of the CCD Data (accuracy of the energy scale etc) is significantly affected by the processing




Contamination on the XIS OBF until 2007Apr evaluated from E0102 obs. (ver.1.x products)



Suzaku/XIS contamination history, $N_c/N_0 = 6$ assumed Ver.2




XIS0, XIS3 spectra of E0102




RXJ1856 observed on 2007Oct

RXJ1856 2007Oct XIS0 XIS3




XIS1(BI) rxj1856_x1_fixnorm.qdp RXJ1856 XIS1 rev2+HEASOFT6.4

RJX1856




Factor of 2 Underestimate
Below 0.3keV



Contamination N_O vs N_C



PKS2155 2005Dec/2006May



*)Error bars are 90% error for 2parameters confidence,i.e., $\Delta\chi^2=4.61$

Factor of 2 underestimate below 0.3keV

- Unable to improve the fit only with C & O
- Absorption by Heavier Element **No apparent edge found**
- **Absorption by H** (or He) but **too much $\sim 10^{21} \text{cm}^2$**
- Constant Factor (Grating Problem at low energy?)




PKS2155 2007Apr

Power * Abs

Index Γ and N_H free & common

Normalization free & vary independently



Users may encounter this level of inconsistency among sensors

Atmospheric Fluorescence Line

- When the telescope is looking at the shining Earth or its atmosphere, fluorescence lines of the Earth atmosphere (N-K, O-K) by Solar X-rays are contaminated in the observed spectra.
- Intensity and line ratio depends on the elevation angle from the Earth rim and the Solar activity.

N-K
(0.39keV)

O-K
(0.52keV)


DAY EARTH

$0 < \text{DYE_ELV} < 5$


$5 < \text{DYE_ELV} < 10$

$10 < \text{DYE_ELV} < 20$

$20 < \text{DYE_ELV} < 30$







Detected Intensity of N-K line can be used to measure the contamination thickness







Radial Profile of Contamination from Atmospheric N-K, O-K


$N_C(10^{18}\text{cm}^{-2})$



Radial Profile of Contamination from Atmospheric O-K for FI-CCD



Cut-Off-Rigidity Dependence of the XIS NXB




The COR dependence of the NXB can be used to model the XIS NXB.

- 1) **Sort** the NXB spectra from the Night Earth Database into each COR bin.
- 2) Calculate the appropriate **weights** of each COR bin for a given source observation.
- 3) **Make** the NXB spectrum for the given source observation by summing the sorted spectra with appropriate weights.

Other parameters, such as PINUD of the HXD, can be used instead of the COR.

5-12keV NXB (integrated for 5ks)

Tawa et al., 2008
PASJ in press



Reproducibility 5-12keV XIS NXB

- 2005Oct-2006Jun (Cut 2005Sep)
- Applying “Orbit Filter”
- 5ks integration
 - Duration >> 5ks

We introduced a new COR database



Parameter	COR	COR2	PINUD
XIS0	4.4±0.5%	3.5±0.7%	2.8±0.9%
XIS1	6.6±0.1%	6.9±0.1%	4.4±0.1%
XIS2	5.7±0.4%	5.7±0.4%	4.0±0.6%
XIS3	3.3±0.9%	2.4±1.2%	3.8±0.8%

% of the NXB

Case Study

- Target is (excess) diffuse emission over XIS FOV with comparable surface density as the average CXB
- Exposure 100ks
- For 5-12keV band integrated counts for the whole area of an FI-CCD
 - CXB: $9.7 \times 10^{-3} \text{ c/s}$
 - Target Diffuse Emission: $9.7 \times 10^{-3} \text{ c/s}$ (assumed)
 - NXB: $52 \times 10^{-3} \text{ c/s}$

• Statistical Error

$$\Delta I_{\text{stat}} = 0.84 \times 10^{-3} \text{ c/s}$$

• Fluctuation of the CXB (estimated) $\Delta I_{\text{CXB}} = 1.5 \times 10^{-3} \text{ c/s}$

• NXB reproducibility (PINUD)

$$\Delta I_{\text{NXB}} = 1.8 \times 10^{-3} \text{ c/s}$$

Comparable

*) CXB fluctuation dominates when we take 1-7keV band.

xisnxbgen is introduced HEASOFT6.4

to make an appropriate NXB spectra for a given source observation

We (XIS team) had been providing perl scripts and the NXB database independent to the HEASOFT/CALDB.

The new ftool **xisnxbgen** is simpler to use, and the NXB data base is now included in the CALDB.

← We have to update every half a year at least

See Detail by fhelp

1. Create an estimated NXB file 'xis1_nxb.pi' in the DET coordinates region 'ds9.reg'

```
% xisnxbgen xis1_nxb.pi xis1_src.pi DETREG ds9.reg suzaku.orb none
```

2. Create an estimated NXB file 'xis1_nxb.pi' in the SKY coordinates region 'ds9.reg' including the calibration source area, and an image in the energy range of $274 \leq PI \leq 548$ (1-2 keV).

```
% xisnxbgen xis1_nxb.pi xis1_src.pi SKYREG ds9.reg  
suzaku.orb suzaku.att pixq_and=0 pi_min=274 pi_max=548
```